

Report for the period October 1, 1993 - September 30, 1994

submitted to:

The Chesapeake Bay Stock Assessment Committee:
attention: M. Elizabeth Gillelan, Division Chief
NOAA Chesapeake Bay Office
National Marine Fisheries Service
410 Severn Avenue, Suite 107A
Annapolis MD 21403

by

The School of Marine Science and Virginia Institute of Marine Science
The College of William and Mary
Gloucester Point, VA 23062
and
Virginia Marine Resources Commission
P.O. Box 756
Newport News, VA 23607-0756

for the program entitled:

**Fishery independent standing stock surveys of oyster populations in the
Virginia sub estuaries of the Chesapeake Bay and a comparison
with continuing estimates obtained from fishery dependent data**

Investigators: Dr. Roger Mann (SMS/VIMS) and Dr. James Wesson (VMRC).

date of report submission: November 4, 1994

Table of Contents

Introduction

History of the Virginia oyster resource and the need for stock assessment

Fishery Independent Sampling

Methods:

Selection of sample numbers and locations (with two figures)

Sampling gear

Data collection

Data reduction and archiving

Results and Discussion:

Data analysis (with 21 figures and 19 statistical summaries)

General summary of population sizes (with one table)

Sizedistribution data and implications for interpretation of general summary (with 2 figures)

Fishery Dependent Methods

(with 1 figure and 3 tables)

Conclusions and recommendations

Literature cited

Acknowledgements: We thank John Register and Calvin Wilson of the Virginia Marine Resources Commission, Reinaldo Morales Alamo and Kenneth Walker of the Virginia Institute of Marine Science, and numerous students of the School of Marine Science, Virginia Institute of Marine Science for their assistance in the field; Gerry Showalter of the Virginia Marine Resources Commission for assistance with reef mapping; and Chris Bonzek and Robert Harris of the Fisheries Data Management Unit of the Department of Fisheries Science at the Virginia Institute of Marine Science of development of the custom database for field data.

Introduction

History of the Virginia oyster resource and the need for stock assessment

Extensive description of the Virginia oyster resource and history of its utilization has been given by Haven, Hargis and Kendall (1981), and more recently reviewed by Hargis and Haven (1988). These contributions, among many others, describe a state of continuing decline. To facilitate resource management a fishery independent survey was proposed to and subsequently supported by the Chesapeake Bay Stock Assessment Committee in 1993. This report covers activity on that program for the period October of 1993 through September of 1994.

Spatial variability in distribution of oysters within an oyster reef system, and distribution of reefs in the intertidal and/or subtidal regions complicate fishery independent estimation of standing stock. By contrast, fishery dependent estimates of oyster standing stock can be made, where adequate data on effort and temporal changes in landings exist, through application of Leslie-DeLury regression analysis (Barber and Mann, 1991). Intensive, fishery independent estimates are rare but pivotal to examination of spawning capabilities of broodstock supporting commercial fisheries and related requirements for establishment of fishery catch quotas. The James River, Virginia has served as the focal point for the Virginia oyster industry for over a century, being the source of the majority of seed oysters that were transplanted for grow-out to locations within the Virginia portion of the Chesapeake Bay and much further afield in the Middle Atlantic states (Haven et al, 1981). The Rappahannock River in Virginia was, for many years, a source of large and valued oysters for both the shucking and half shell trade. It is surprising that comparatively little effort has been previously expended to estimate standing stock in both the James and Rappahannock Rivers given the acknowledged need for such data in fishery management. Continuing losses of productive oyster reef over the past three decades to Haplosporidium nelsoni, commonly known as MSX, and Perkinsus marinus, commonly known as "Dermo", in the higher salinity regions of both rivers, combined with increased fishing pressure on all remaining stocks, have emphasized the need for working estimates of standing stock. This need has been further exaggerated in the James River by a change in emphasis in the past decade from the harvesting of "seed" oysters to larger "market" oysters, and the reduction in size limit of the latter from three to two-and-one-half inches maximum dimension (although this action was reversed with an increase in minimum market size to three inches for the 1994-1995 season). The fishery is now facing the dilemma of exploiting the limited remaining broodstock from the James River in order to retain a viable fishery for "market" oysters, while simultaneously threatening the long term future of the river as a seed producing location.

Fishery Independent Sampling

The primary objective of the study was to effect a fishery independent study of the standing stock of oysters, both market and seed, in the Virginia portion of the Chesapeake Bay and the Seaside of the Eastern Shore. For the period reported here the focus of activity was on the James and Rappahannock Rivers.

Methods

The selection of sample numbers and locations

James River

The initial focus of the program was the oyster resource of the James River. We designed a quantitative sampling program using quadrats located in a random grid placed over a map of the known oyster resources in the James. In essence, this is a stratified random grid with the documented oyster reefs or rocks forming the strata. The area surveyed is described in extensive surveys made by VIMS and reported by Haven and Whitcomb (1983). These areas have been subjected to regular survey by VMRC and VIMS personnel for at least two decades by dredge. The limits of the known oyster reef were mapped by the Surveying Engineering Department at VMRC and the grids for sampling set with Loran coordinates (Loran was checked daily when in the field from known markers at both the beginning and end of the day). Sampling areas are described in Figures 1 and 2. Figure 1 relates sampling areas to bottom type. Figure 2 identifies the sampled rocks by number. These numbers are used throughout this report in summary tables and graphics. Sampling areas 1 through 11 in Figure 2 represent the limits of hard oyster rock strata selected, mapped and sampled within the larger public oyster grounds in those regions. The limits of hard oyster rock strata within sampling areas 12 through 19 were not mapped separately because of the large areas involved; consequently, we knew beforehand that sampling grids selected in areas 12-19 would include both oyster rock strata as well as bare sandy or muddy strata. Sampling sites were picked by random numbers within the grids and oysters were sampled with a hydraulically operated patent tong. In this manner a total of 823 stations were occupied in the James River.

Rappahannock River


The sampling protocol for the Rappahannock River was as for the James River and employed a quantitative sampling program using quadrats located in a random grid placed over a map of the known oyster resources. Although once extensive, these are now limited to the upper part of the Rappahannock above Bowlers Rock and Morattico Bar. The only commercially exploited reef of any consequence is Russ' Rock. The reefs were again the basis for stratified random sampling. The area surveyed is described in Haven and Whitcomb (1989). The limits of the known oyster reef were mapped by the Surveying Engineering Department at VMRC and the grids for sampling set with Loran coordinates. Loran was, again, checked daily when in the field from known markers at both the beginning and end of the day. Sampling sites were picked by random numbers within the grids and oysters were sampled with a hydraulically operated patent tong.

Sampling gear

Both tongs and dredges are commonly used to examine oyster populations; however, only the former are good quantitative tools (see Chai et al, 1992). Initially, we examined a standard patent tong of known area; however, tests proved this to be an unpredictable sampling tool in that penetration into the hard bottom on the reef surface was inconsistent resulting in high variability in replicate samples on the same site. We replaced the tong with an hydraulically operated tong which separates the closing actions of the tong from the retrieval action. This has proven to be vastly superior in providing consistent penetration of

James River Public (Baylor) Oyster Grounds

Figure 1




Outline of areas sampled during Fall 1993 oyster stock assessment survey: superimposed over chart of bottom types modified from Haven, Whitcomb and Kendall (1981) by the VIMS Center for Coastal Management and Policy. Areas in white represent soft mud primarily.

Figure 2

James River Oyster Stock Assessment: Fall 1993

Location of oyster rocks sampled. Individual rocks are as in the previous description of bottom type and are identified by number in the key. These numbers cross reference with the summary table of standing stock estimates, all data plots relating to sampling adequacy, and data summary tables for the individual rocks.



the bottom and replication sampling. The hydraulic tong was installed on the VMRC vessel Wolftrap. This vessel was used in all survey work described herein.

Data collection

The open dimensions of the tong were such that it sampled one square meter. Upon retrieval the sample was washed on the cull board and processed for counts of live oysters as spat (young of the year), small oysters (less than 2.5 inches), and market (greater than 2.5 inches) oysters. In addition, the opportunity was taken to collect data on dead oysters with paired valves (boxes, indicating recent mortality). The volume of shell retrieved in each tong was also recorded as an index of the quantity of cultch material present at each station. Between six and nine people were on board on each day of sampling, and all were trained to avoid inconsistency in categorization of oysters. This process was much more labor intensive than originally envisaged, with between 30 and 60 samples being processed each day depending on weather conditions, crew size and the time required to wash and separate samples. Sampling of the James River was completed in late December of 1993. A further effort in early January of 1994 focused on sampling in the Rappahannock River.

Data reduction and archiving

A custom database program for field data was developed by the Fisheries Data Management Unit in the Department of Fisheries Science at the School of Marine Science and Virginia Institute of Marine Science. Archived material is available on request. Size distribution data was additionally archived and preliminary analysis effected using commercial spreadsheet software (Microsoft Excel).

Results and Discussion

Data analysis

Prior to using data to estimate standing stock initial questions relating to sampling design and adequacy need to be addressed. As mentioned earlier, there is a lack of previous quantitative assessment data for this resource. Two primary questions arise:

1. Are there strata reasonable? The background behind this question is that recent surveys by Haven and Whitcomb (1983, 1989) illustrate varying bottom type within the chosen strata - from mud to hard shell bottom. This could present a significant sampling problem in that strata are sufficiently heterogeneous to be of limited ecological and statistical value.
2. Assuming 1 (above) is not a problem, are there sufficient samples to adequately represent the strata and allow estimates of abundance per unit area and, subsequently, total standing stock.

Bros and Cowell (1987) offer a good discussion of methods of estimating sample size in situations where minimum detectable difference cannot be specified a priori, as is the case in this situation. Their proposed method incorporates use of resolving power as a primary factor and sampling feasibility (an issue here with time and cost) as a secondary factor. They suggest the standard error of the mean be used as a measure of appropriate sampling effort. We have adopted their suggestion.

Questions 1 and 2 above were primarily addressed by a single analysis in which data were examined collectively within each strata. A plot was generated of mean number of oysters per patent tong (one square meter) sample and standard error of the mean versus number of samples included in the calculation. This calculation was repeated ten times for data within a strata with samples being chosen at random from those available. Random sampling eliminated any bias that resulted from sequential data entry in accordance with sampling in the field sampling (the latter may have resulted, inadvertently in temporally focused sampling on a particular substrate type). In a regime where variability with bottom type was high and the sample size was low then the mean would not stabilize, and where sampling was insufficient the standard error of the mean would not demonstrate a stable trend of decreasing value - remembering of course that the standard error value will eventually continue to decrease with increasing number of samples included in the calculation because the standard error is inversely proportional to the square root of the number of observations of the mean. Increasing sample size will eventually solve both these problems, but the number of samples required might be very large.

Figures 3 through 23 provide the visual description of all 19 sampling areas in the James when subjected to plotting estimates of the standard error of the mean versus number of samples collected (included in the analysis) for each of the sampling areas. The bold numeral in the bottom left corner of each plot provides the cross reference to Figure 2. As mentioned earlier, sampling areas 1 through 11 (figures 3 through 15) represent the limits of well defined hard oyster rock strata within the public oyster grounds in those regions. Sampling grids selected in areas 12-19 (figures 16 through 23) included both oyster rock strata as well as bare sandy or muddy strata, and heterogeneity within the strata was expected to be more acute in these areas. The area of each reef was quite variable, and resulted in a variation in the number of stations sampled. Each figure represents the means of 10 randomized groups of samples subjected to the previously described analysis for each strata. In addition, areas 4 (Horsehead Middle) and 8 (Point of Shoals) have plots generated with one randomized set of data for the strata. These pairs of plots are identified with the bold characters 4 and 4b, and 8 and 8b respectively (figures 6 and 7, and 11 and 12 respectively). There is generally good agreement between the plots (figure 6 versus 7, and 11 versus 12). These plots suggest adequate sampling within the strata to account for bottom type variability and for general spatial coverage.

Interspersed between the individual plots of standard error versus number of samples included are statistical summaries for the corresponding sampling areas. Upper Deep Water Shoal (Figure 3) is a large area (234 acres) and the mean did not stabilize until approximately 40 samples were included in the analysis, although the standard error measurements settled into a steady decline at about 30 samples. This is a good example of an area with small scale bottom variability directly adjacent to a deep channel. Future surveys will not require a repetition of 99 individual samples. The standard error remains high at approximately 10 even with 60 samples included in the analysis. This is indicative of limited but consistent small scale heterogeneity in abundance within the strata; however, our sampling was more than adequate to compensate for this. Lower Deep Water Shoal (Figure 4) is much smaller by comparison (20 acres) but gave relatively stable estimates of mean oyster density at small sample numbers.

Some reefs such as Upper Horsehead (Figure 5, 3 acres), Moon Rock (Figure 9, 4 acres), Shanty Rock (Figure 14, 3.5 acres), and Dry Lumps (Figure 15, 6 acres), are all rather small with limited sample numbers. Despite this the analysis suggests relative homogeneity within the small strata and good estimates of standing stock.

Some of the larger reefs with well defined hard rock bottom types (areas 1 through 11 on Figure 2, corresponding to figures 3 through 15) such as Horsehead middle and lower, V Rock, Point of Shoals, and Cross Rock all exhibit stability in estimates of the mean with fairly small sample sizes. This is also the case for some of the strata with less well defined bottoms such as Swash & Swash Mud Slough (Figure 19). This suggests, again, that future studies will require a smaller number of sample stations than were employed in 1993. Moving to strata with less well defined bottom types (areas 12 through 19 on Figure 2, corresponding to Figures 16 through 23) was often accompanied by a decrease in mean oyster density (compare Mulberry Point at 24 per sample (Figure 16) and Swash at 4.8 per sample (Figure 17) with earlier values for Deep Water Shoal, Point of Shoals, and V Rock). Care is required when examining some of these latter plots (for example Figures 21 and 22, Jail Island Lower and Offshore) in that the values appear initially unstable up to quite large sample sizes - this is a function of the axes in that the absolute numbers are small by comparison with earlier plots. Jail Island Offshore in particular is a very large area with low oyster density. Wreck Shoal (Figure 23) is the most down stream position of all of the sampled strata and has been subjected to intense disease pressure for the past several years with corresponding mortalities. It is also a very large reef (585 acres). Both the mean and the standard error values stabilize when between 40 and 50 samples included in the analysis. The abundance data of less than 9 oysters per square meter illustrate the cumulative disease impact in that this was a major oyster producing reef in the upper James during the 1982-1986 period when market oyster production was high.

James River Patent Tong Survey

Comparison of Sample Size to Standard Error & Mean

Deep Water Shoal - Upper




Figure 3


STATISTICAL SUMMARY FOR UPPER DEEP WATER SHOAL (1)

	LIVE OYSTERS				BOXES				SHELL VOL (I)
	SPAT	SMALL	MARKET	TOTAL SML+MKT	SPAT	SML	MKT	TOTAL SML+MKT	
LOW VALUE (PER SQ. METER)	0	0	0	0		0	0	0	0
HIGH VALUE (PER SQ. METER)	237	84	84	291		16	22	22	20
MEAN NO. PER SQ. METER	49.1	19.8	19.8	68.9		1.7	0.9	2.6	5.6
STD.DEV.	60.4	23.8	23.8	78.3		2.6	2.7	3.8	5.5
NO. SAMPLES (n)	99	99	99	99		99	99	99	99
STD. ERROR (SE)	6.1	2.4	2.4	7.9		0.3	0.3	0.4	0.6
t.05 VALUE FOR n (t.05)	2.571	2.571	2.571	2.571		2.571	2.571	2.571	2.571
(SE)*(t.05) = t.05SE	15.61	6.16	6.16	20.23		0.67	0.69	0.99	1.42
UPPER 95% CONF. INTVL. FOR MEAN	64.7	26.0	26.0	89.2		2.3	1.6	3.6	7.0
LOWER 95% CONF. INTVL FOR MEAN	33.5	13.7	13.7	48.7		1.0	0.2	1.6	4.2
CONVERS. FACTOR ACRES TO SQ. METERS	4046.9	4046.9	4046.9	4046.9					
ESTIMATED NO. OYSTERS IN 1 BUSHEL	1000	500	500	500					
ROCK ACREAGE	233.92	233.92	233.92	233.92					
MEAN NO. BUSHEL PER ACRE	198.7	160.6	160.6	359					
UPPER 95% CONF. INTVL. FOR THIS MEAN	261.8	210.5	210.5	472					
LOWER 95% CONF. INTVL. FOR THIS MEAN	135.5	110.8	110.8	246					
MEAN TOTAL BUSHEL IN WHOLE ROCK	46472	37579	37579	84051					
UPPER 95% CONF. INTVL. FOR THIS MEAN	61252	49240	49240	110492					
LOWER 95% CONF. INTVL. FOR THIS MEAN	31692	25918	25918	57610					

James River Patent Tong Survey

Comparison of Sample Size to Standard Error & Mean

Deep Water Shoal - Lower



Number of Samples Included

STATISTICAL SUMMARY FOR LOWER DEEP WATER SHOAL (2)

	LIVE OYSTERS				BOXES				SHELL VOL (I)
	SPAT	SMALL	MARKET	TOTAL SML+MKT	SPAT	SML	MKT	TOTAL SML+MKT	
LOW VALUE (PER SQ. METER)	0	0	0	0		0	0	0	0.5
HIGH VALUE (PER SQ. METER)	28	21	21	44		6	8	12	17
MEAN NO. PER SQ. METER	9.9	8.5	8.5	18.4		1.3	1.1	2.4	5.9
STD.DEV.	7.7	5.3	5.3	12.5		1.5	2.1	3.2	4.4
NO. SAMPLES (n)	20	20	20	20		20	20	20	20
STD. ERROR (SE)	1.7	1.2	1.2	2.8		0.3	0.5	0.7	1.0
t.05 VALUE FOR n (t.05)	2.093	2.093	2.093	2.093		2.093	2.093	2.093	2.093
(SE)*(t.05) = t.05SE	3.60	2.49	2.49	5.86		0.71	0.99	1.49	2.07
UPPER 95% CONF. INTVL. FOR MEAN	13.5	11.0	11.0	24.3		2.0	2.1	3.8	7.9
LOWER 95% CONF. INTVL FOR MEAN	6.3	6.0	6.0	12.5		0.5	0.1	0.9	3.8
CONVERS. FACTOR ACRES TO SQ. METERS	4046.9	4046.9	4046.9	4046.9					
ESTIMATED NO. OYSTERS IN 1 BUSHEL	1000	500	500	500					
ROCK ACREAGE	19.93	19.93	19.93	19.93					
MEAN NO. BUSHEL PER ACRE	40	69	69	109					
UPPER 95% CONF. INTVL. FOR THIS MEAN	55	89	89	144					
LOWER 95% CONF. INTVL. FOR THIS MEAN	26	49	49	74					
MEAN TOTAL BUSHEL IN WHOLE ROCK	798	1371	1371	2170					
UPPER 95% CONF. INTVL. FOR THIS MEAN	1089	1773	1773	2862					
LOWER 95% CONF. INTVL. FOR THIS MEAN	508	969	969	1477					

James River Patent Tong Survey

Comparison of Sample Size to Standard Error & Mean

Horsehead - Upper




Figure 5


STATISTICAL SUMMARY FOR UPPER HORSEHEAD (3)

	LIVE OYSTERS				BOXES				SHELL VOL. (I)
	SPAT	SMALL	MARKET	TOTAL SML+MKT	SPAT	SML	MKT	TOTAL SML+MKT	
LOW VALUE (PER SQ. METER)		179	36	232		7	0	7	1.5
HIGH VALUE (PER SQ. METER)		382	77	459		33	2	33	10
MEAN NO. PER SQ. METER		294.7	55.3	350.0		17.7	0.7	18.3	7.2
STD.DEV.		104.4	20.6	113.8		13.6	1.2	13.3	4.9
NO. SAMPLES (n)		3	3	3		3	3	3	3
STD. ERROR (SE)		60.3	11.9	65.7		7.9	0.7	7.7	2.8
t.05 VALUE FOR n (t.05)		4.303	4.303	4.303		2.093	2.093	2.093	2.093
(SE)*(t.05) = t.05SE		259.42	51.18	282.64		16.45	1.40	16.09	5.93
UPPER 95% CONF. INTVL. FOR MEAN		554.1	106.5	632.6		34.1	2.1	34.4	13.1
LOWER 95% CONF. INTVL FOR MEAN		35.2	4.2	67.4		1.2	-0.7	2.2	1.2
CONVERS. FACTOR ACRES TO SQ. METERS		4046.9	4046.9						
ESTIMATED NO. OYSTERS IN 1 BUSHEL		1000	500						
ROCK ACREAGE		3.009	3.009						
MEAN NO. BUSHEL PER ACRE		1192	448	1640					
UPPER 95% CONF. INTVL. FOR THIS MEAN		2242	862	3104					
LOWER 95% CONF. INTVL. FOR THIS MEAN		143	34	176					
MEAN TOTAL BUSHEL IN WHOLE ROCK		3588	1348	4936					
UPPER 95% CONF. INTVL. FOR THIS MEAN		6747	2594	9341					
LOWER 95% CONF. INTVL. FOR THIS MEAN		429	101	530					

James River Patent Tong Survey

Comparison of Sample Size to Standard Error & Mean


Horsehead - Middle



James River Patent Tong Survey

Comparison of Sample Size to Standard Error & Mean

Horsehead - Middle



STATISTICAL SUMMARY FOR MIDDLE HORSEHEAD (4)

	LIVE OYSTERS				BOXES				SHELL VOL (0)
	SPAT	SMALL	MARKET	TOTAL SML+MKT	SPAT	SML	MKT	TOTAL SML+MKT	
LOW VALUE (PER SQ. METER)	64	0	65		4	0	5	2	
HIGH VALUE (PER SQ. METER)	579	32	594		32	2	32	26	
MEAN NO. PER SQ. METER	214.3	7.4	221.6		13.5	0.4	13.9	13.0	
STD.DEV.	138.7	8.5	141.2		8.8	0.6	8.7	6.6	
NO. SAMPLES (n)	20	20	20		20	20	20	20	
STD. ERROR (SE)	31.0	1.9	31.6		2.0	0.1	1.9	1.5	
t.05 VALUE FOR n (t.05)	2.093	2.093	2.093		2.093	2.093	2.093	2.093	
(SE)*(t.05) = t.05SE	64.91	3.97	66.07		4.11	0.28	4.08	3.10	
UPPER 95% CONF. INTVL. FOR MEAN	279.2	11.3	287.7		17.6	0.7	17.9	16.1	
LOWER 95% CONF. INTVL. FOR MEAN	149.3	3.4	155.5		9.3	0.1	9.8	9.9	
CONVERS. FACTOR ACRES TO SQ. METERS	4046.9	4046.9							
ESTIMATED NO. OYSTERS IN 1 BUSHEL	1000	500							
ROCK ACREAGE	19.465	19.465							
MEAN NO. BUSHEL PER ACRE	867	59	927						
UPPER 95% CONF. INTVL. FOR THIS MEAN	1130	92	1221						
LOWER 95% CONF. INTVL. FOR THIS MEAN	604	27	632						
MEAN TOTAL BUSHEL IN WHOLE ROCK	16877	1158	18035						
UPPER 95% CONF. INTVL. FOR THIS MEAN	21990	1784	23774						
LOWER 95% CONF. INTVL. FOR THIS MEAN	11764	532	12297						

James River Patent Tong Survey

Comparison of Sample Size to Standard Error & Mean

Horsehead - Lower




Figure 8

STATISTICAL SUMMARY FOR LOWER HORSEHEAD (5)

	LIVE OYSTERS				BOXES				SHELL VOL (I)
	SPAT	SMALL	MARKET	TOTAL SML+MKT	SPAT	SML	MKT	TOTAL SML+MKT	
LOW VALUE (PER SQ. METER)	15	0	0	15		4	0	6	6
HIGH VALUE (PER SQ. METER)	470	110	110	484		19	4	19	22
MEAN NO. PER SQ. METER	253.4	18.8	18.8	272.2		9.8	0.9	10.6	13.1
STD.DEV.	120.5	23.6	23.6	125.0		4.3	1.3	3.7	4.3
NO. SAMPLES (n)	20	20	20	20		20	20	20	20
STD. ERROR (SE)	26.9	5.3	5.3	28.0		1.0	0.3	0.8	1.0
t.05 VALUE FOR n (t.05)	2.093	2.093	2.093	2.093		2.093	2.093	2.093	2.093
(SE)*(t.05) = t.05SE	56.40	11.06	11.06	58.52		2.00	0.59	1.73	2.00
UPPER 95% CONF. INTVL. FOR MEAN	309.8	29.8	29.8	330.7		11.8	1.4	12.3	15.1
LOWER 95% CONF. INTVL. FOR MEAN	197.0	7.7	7.7	213.6		7.7	0.3	8.9	11.1
CONVERS. FACTOR ACRES TO SQ. METERS	4046.9	4046.9	4046.9	4046.9					
ESTIMATED NO. OYSTERS IN 1 BUSHEL	1000	500	500	500					
ROCK ACREAGE	19.467	19.467	19.467	19.467					
MEAN NO. BUSHELS PER ACRE	1025	152	152	1177					
UPPER 95% CONF. INTVL. FOR THIS MEAN	1254	241	241	1495					
LOWER 95% CONF. INTVL. FOR THIS MEAN	797	62	62	859					
MEAN TOTAL BUSHELS IN WHOLE ROCK	19963	2954	2954	22917					
UPPER 95% CONF. INTVL. FOR THIS MEAN	24406	4697	4697	29104					
LOWER 95% CONF. INTVL. FOR THIS MEAN	15520	1211	1211	16731					

James River Patent Tong Survey

Comparison of Sample Size to Standard Error & Mean




Figure 9

STATISTICAL SUMMARY FOR MOON ROCK (6)

	LIVE OYSTERS				BOXES				SHELL VOL (I)
	SPAT	SMALL	MARKET	TOTAL	SPAT	SML	MKT	TOTAL	
			SML+MKT				SML+MKT		
LOW VALUE (PER SQ. METER)	148	10	158		3	0	3		8
HIGH VALUE (PER SQ. METER)	338	36	368		16	2	18		18
MEAN NO. PER SQ. METER	247.0	24.8	271.8		10.5	0.5	11.0		14.0
STD. DEV.	86.7	11.2	96.7		5.4	1.0	6.2		4.5
NO. SAMPLES (n)	4	4	4		4	4	4		4
STD. ERROR (SE)	43.3	5.6	48.4		2.7	0.5	3.1		2.3
t.05 VALUE FOR n (t.05)	3.182	3.182	3.182		3.182	3.182	3.182		3.182
(SE)*(t.05) = t.05SE	137.88	17.78	153.93		8.67	1.59	9.81		7.23
UPPER 95% CONF. INTVL. FOR MEAN	384.9	42.5	425.7		19.2	2.1	20.8		21.2
LOWER 95% CONF. INTVL. FOR MEAN	109.1	7.0	117.8		1.8	-1.1	1.2		6.8
CONVERS. FACTOR ACRES TO SQ. METERS	4046.9	4046.9							
ESTIMATED NO. OYSTERS IN 1 BUSHEL	1000	500							
ROCK ACREAGE	3.95	3.95							
MEAN NO. BUSHEL PER ACRE	999.6	200.3	1200						
UPPER 95% CONF. INTVL. FOR THIS MEAN	1557.6	344.2	1902						
LOWER 95% CONF. INTVL. FOR THIS MEAN	441.6	56.4	498						
MEAN TOTAL BUSHEL IN WHOLE ROCK	3948	791	4740						
UPPER 95% CONF. INTVL. FOR THIS MEAN	6152	1360	7512						
LOWER 95% CONF. INTVL. FOR THIS MEAN	1744	223	1967						

James River Patent Tong Survey

Comparison of Sample Size to Standard Error & Mean




Figure 10

STATISTICAL SUMMARY FOR V-ROCK (7)

	LIVE OYSTERS				BOXES				SHELL VOL (l)
	SPAT	SMALL	MARKET	TOTAL SML+MKT	SPAT	SML	MKT	TOTAL SML+MKT	
LOW VALUE (PER SQ. METER)	0	0	0	0		0	0	0	0
HIGH VALUE (PER SQ. METER)	99	326	43	344		30	7	37	25
MEAN NO. PER SQ. METER	30.9	157.6	20.3	173.4		10.8	1.2	12.0	14.4
STD.DEV.	38.4	62.6	10.5	74.7		6.7	1.6	8.3	4.6
NO. SAMPLES (n)	15	36	36	36		36	36	36	36
STD. ERROR (SE)	9.9	10.4	1.7	12.4		1.1	0.3	1.4	0.8
t.05 VALUE FOR n (t.05)	2.131	2.029	2.029	2.029		2.029	2.029	2.029	2.029
(SE)*(t.05) = t.05SE	21.125	21.1589	3.5401	25.25119		2.253	0.537	2.789946	1.568419
UPPER 95% CONF. INTVL. FOR MEAN	52.1	178.7	23.8	198.7		13.1	1.8	14.8	16.0
LOWER 95% CONF. INTVL. FOR MEAN	9.8	136.4	16.8	148.2		8.6	0.7	9.2	12.9
CONVERS. FACTOR ACRES TO SQ. METERS		4046.9	4046.9						
ESTIMATED NO. OYSTERS IN 1 BUSHEL		1000	500						
ROCK ACREAGE		72.053	72.053						
MEAN NO. BUSHEL PER ACRE		638	164	802					
UPPER 95% CONF. INTVL. FOR THIS MEAN		723	193	916					
LOWER 95% CONF. INTVL. FOR THIS MEAN		552	136	688					
MEAN TOTAL BUSHEL IN WHOLE ROCK		45950	11842	57792					
UPPER 95% CONF. INTVL. FOR THIS MEAN		52120	13906	66026					
LOWER 95% CONF. INTVL. FOR THIS MEAN		39780	9777	49557					

James River Patent Tong Survey

Comparison of Sample Size to Standard Error & Mean




Figure 11

Mean Oysters per Sample


8

Number of Samples Included

James River Patent Tong Survey

Comparison of Sample Size to Standard Error & Mean

Point of Shoals



STATISTICAL SUMMARY FOR POINT OF SHOALS (8)

	LIVE OYSTERS				BOXES				SHELL VOL (I)
	SPAT	SMALL	MARKET	TOTAL SML+MKT	SPAT	SML	MKT	TOTAL SML+MKT	
LOW VALUE (PER SQ. METER)	3	2	0	0		0	0	0	0.5
HIGH VALUE (PER SQ. METER)	253	69	299		28	6	28	17	17
MEAN NO. PER SQ. METER	104.9	23.9	128.6		5.1	1.5	6.6	8.0	8.0
STD.DEV.	72.7	17.2	87.9		5.7	1.5	5.8	4.6	4.6
NO. SAMPLES (n)	35	35	35		35	35	35	35	35
STD. ERROR (SE)	12.3	2.9	14.9		1.0	0.3	1.0	0.8	0.8
t.05 VALUE FOR n (t.05)	1.982	1.982	1.982		1.982	1.982	1.982	1.982	1.982
(SE)*(t.05) = t.05SE	24.36	5.77	29.45		1.92	0.50	1.94	1.55	1.55
UPPER 95% CONF. INTVL. FOR MEAN	129.3	29.7	158.1		7.1	2.0	8.5	9.5	9.5
LOWER 95% CONF. INTVL. FOR MEAN	80.5	18.1	99.2		3.2	1.0	4.7	6.4	6.4
CONVERS. FACTOR ACRES TO SQ. METERS	4046.9	4046.9							
ESTIMATED NO. OYSTERS IN 1 BUSHEL	1000	500							
ROCK ACREAGE	131.71	131.71							
MEAN NO. BUSHEL PER ACRE	424	193	618						
UPPER 95% CONF. INTVL. FOR THIS MEAN	523	240	763						
LOWER 95% CONF. INTVL. FOR THIS MEAN	326	147	472						
MEAN TOTAL BUSHEL IN WHOLE ROCK	55906	25463	81369						
UPPER 95% CONF. INTVL. FOR THIS MEAN	68893	31617	100509						
LOWER 95% CONF. INTVL. FOR THIS MEAN	42919	19309	62229						

James River Patent Tong Survey

Comparison of Sample Size to Standard Error & Mean




Figure 13

STATISTICAL SUMMARY FOR CROSS ROCK (9)

	LIVE OYSTERS				BOXES				SHELL VOL (I)
	SPAT	SMALL	MARKET	TOTAL SML+MKT	SPAT	SML	MKT	TOTAL SML+MKT	
LOW VALUE (PER SQ. METER)	3	1		6		0		0	0.5
HIGH VALUE (PER SQ. METER)	193	18		204		39	8	47	23
MEAN NO. PER SQ. METER	75.1	7.8		82.9		12.7	2.9	15.6	13.0
STD.DEV.	57.5	4.5		60.6		10.7	2.3	12.2	7.4
NO. SAMPLES (n)	19	19		19		19	19	19	19
STD. ERROR (SE)	13.2	1.0		13.9		2.5	0.5	2.8	1.7
t.05 VALUE FOR n (t.05)	2.093	2.093		2.093		2.093	2.093	2.093	2.093
(SE)*(t.05) = t.05SE	27.62	2.17		29.10		5.14	1.10	5.84	3.55
UPPER 95% CONF. INTVL. FOR MEAN	102.7	10.0		112.0		17.8	4.0	21.4	16.5
LOWER 95% CONF. INTVL. FOR MEAN	47.5	5.7		53.9		7.5	1.8	9.7	9.4
CONVERS. FACTOR ACRES TO SQ. METERS	4046.9	4046.9							
ESTIMATED NO. OYSTERS IN 1 BUSHEL	1000	500							
ROCK ACREAGE	36.688	36.688							
MEAN NO. BUSHEL PER ACRE	303.9	63.5		367					
UPPER 95% CONF. INTVL. FOR THIS MEAN	415.7	81.0		497					
LOWER 95% CONF. INTVL. FOR THIS MEAN	192.2	45.9		238					
MEAN TOTAL BUSHEL IN WHOLE ROCK	11151	2329		13480					
UPPER 95% CONF. INTVL. FOR THIS MEAN	15252	2972		18224					
LOWER 95% CONF. INTVL. FOR THIS MEAN	7050	1685		8735					

James River Patent Tong Survey

Comparison of Sample Size to Standard Error & Mean

Shanty Rock




Figure 14

STATISTICAL SUMMARY FOR SHANTY ROCK (10)

	LIVE OYSTERS				BOXES				SHELL VOL (I)
	SPAT	SMALL	MARKET	TOTAL SML+MKT	SPAT	SML	MKT	TOTAL SML+MKT	
LOW VALUE (PER SQ. METER)	7	0	0	7		2	0	4	10
HIGH VALUE (PER SQ. METER)	48	5	5	51		29	7	36	14
MEAN NO. PER SQ. METER	32.5	2.3	2.3	34.8		14.3	3.0	17.3	12.8
STD.DEV.	19.0	2.1	2.1	20.7		11.1	2.9	13.5	1.9
NO. SAMPLES (n)	4	4	4	4		4	4	4	4
STD. ERROR (SE)	9.5	1.0	1.0	10.3		5.6	1.5	6.7	0.9
t.05 VALUE FOR n (t.05)	3.182	3.182	3.182	3.182		3.182	3.182	3.182	3.182
(SE)*(t.05) = t.05SE	30.26	3.28	3.28	32.87		17.7	4.68	21.40	3.01
UPPER 95% CONF. INTVL. FOR MEAN	62.8	5.5	5.5	67.6		32.0	7.7	38.6	15.8
LOWER 95% CONF. INTVL. FOR MEAN	2.2	-1.0	-1.0	1.9		-3.5	-1.7	-4.1	9.7
CONVERTS. FACTOR ACRES TO SQ. METERS	4046.9	4046.9	4046.9						
ESTIMATED NO. OYSTERS IN 1 BUSHEL	1000	500	500						
ROCK ACREAGE	3.58	3.58	3.58						
MEAN NO. BUSHELS PER ACRE	132	18	18	150					
UPPER 95% CONF. INTVL. FOR THIS MEAN	254	45	45	299					
LOWER 95% CONF. INTVL. FOR THIS MEAN	9	0	0	9					
MEAN TOTAL BUSHELS IN WHOLE ROCK	471	65	65	536					
UPPER 95% CONF. INTVL. FOR THIS MEAN	909	160	160	1069					
LOWER 95% CONF. INTVL. FOR THIS MEAN	32	0	0	32					

James River Patent Tong Survey

Comparison of Sample Size to Standard Error & Mean




Figure 15

STATISTICAL SUMMARY FOR DRY LUMPS (11)

	LIVE OYSTERS				BOXES				SHELL VOL.(I)
	SPAT	SMALL	MARKET	TOTAL SML+MKT	SPAT	SML	MKT	TOTAL SML+MKT	
LOW VALUE (PER SQ. METER)	3	0	0	3		2	0	2	4
HIGH VALUE (PER SQ. METER)	25	2	2	26		14	2	16	39
MEAN NO. PER SQ. METER	15.0	0.8	0.8	15.8		6.3	0.5	6.8	21.3
STD.DEV.	8.6	1.0	1.0	8.9		4.1	0.8	4.8	13.5
NO. SAMPLES (n)	6	6	6	6		6	6	6	6
STD. ERROR (SE)	3.5	0.4	0.4	3.6		1.7	0.3	1.9	5.5
t.05 VALUE FOR n (t.05)	2.571	2.571	2.571	2.571		2.571	2.571	2.571	2.571
(SE)*(t.05) = t.05SE	9.08	1.03	1.03	9.33		4.29	0.88	4.99	14.17
UPPER 95% CONF. INTVL. FOR MEAN	24.1	1.9	1.9	25.2		10.6	1.4	11.8	35.5
LOWER 95% CONF. INTVL FOR MEAN	5.9	-0.2	-0.2	6.5		2.0	-0.4	1.8	7.2
CONVERS. FACTOR ACRES TO SQ. METERS	4046.9	4046.9	4046.9	4046.9					
ESTIMATED NO. OYSTERS IN 1 BUSHEL	1000	500	500	500					
ROCK ACREAGE	5.93	5.93	5.93	5.93					
MEAN NO. BUSHEL PER ACRE	60.7	6.7	6.7	67					
UPPER 95% CONF. INTVL. FOR THIS MEAN	97.4	15.1	15.1	113					
LOWER 95% CONF. INTVL. FOR THIS MEAN	24.0	0.0	0.0	24					
MEAN TOTAL BUSHEL IN WHOLE ROCK	360	40	40	400					
UPPER 95% CONF. INTVL. FOR THIS MEAN	578	90	90	667					
LOWER 95% CONF. INTVL. FOR THIS MEAN	142	0	0	142					

James River Patent Tong Survey

Comparison of Sample Size to Standard Error & Mean

Mulberry Point




Figure 16

STATISTICAL SUMMARY FOR MULBERRY POINT (12)
 (Combination of "Mulberry Pt." and "Mulberry-Swash")

	LIVE OYSTERS			BOXES				SHELL VOL (I)
	SPAT	SMALL	MARKET	TOTAL SML+MKT	SPAT	SML	MKT	
LOW VALUE (PER SQ. METER)	0	0	0	0		0	0	0
HIGH VALUE (PER SQ. METER)	160	39		199		8	5	11
MEAN NO. PER SQ. METER	18.3	5.6		23.9		1.6	0.6	3.1
STD. DEV.	32.8	8.0		40.3		2.1	1.0	3.0
NO. SAMPLES (n)	45	45		45		45	45	27
STD. ERROR (SE)	4.9	1.2		6.0		0.3	0.1	0.6
t.05 VALUE FOR n (t.05)	2.013	2.013		2.013		2.013	2.013	2.013
(SE)*(t.05) = t.05SE	9.85	2.41		12.10		0.63	0.30	1.17
UPPER 95% CONF. INTVL. FOR MEAN	28.2	8.0		36.0		2.2	0.9	4.2
LOWER 95% CONF. INTVL. FOR MEAN	8.5	3.2		11.8		0.9	0.3	1.9
CONVERS. FACTOR ACRES TO SQ. METERS	4046.9	4046.9		4046.9				
ESTIMATED NO. OYSTERS IN 1 BUSHEL	1000	500		500				
ROCK ACREAGE	86.85	86.85		86.85				
MEAN NO. BUSHEL PER ACRE	74	45		119				
UPPER 95% CONF. INTVL. FOR THIS MEAN	114	65		179				
LOWER 95% CONF. INTVL. FOR THIS MEAN	34	26		60				
MEAN TOTAL BUSHEL IN WHOLE ROCK	6436	3937		10372				
UPPER 95% CONF. INTVL. FOR THIS MEAN	9899	5633		15532				
LOWER 95% CONF. INTVL. FOR THIS MEAN	2973	2240		5213				

James River Patent Tong Survey

Comparison of Sample Size to Standard Error & Mean

Swash




Figure 17


STATISTICAL SUMMARY FOR SWASH (13)

	LIVE OYSTERS				BOXES				SHELL VOL (l)
	SPAT	SMALL	MARKET	TOTAL SML+MKT	SPAT	SML	MKT	TOTAL SML+MKT	
LOW VALUE (PER SQ. METER)	0	0	0	0		0	0	0	0
HIGH VALUE (PER SQ. METER)	19	7	25	25		8	5	9	9
MEAN NO. PER SQ. METER	3.5	1.3	4.8	4.8		0.7	0.5	1.2	1.2
STD.DEV.	5.3	2.0	7.1	7.1		1.9	1.2	2.3	2.3
NO. SAMPLES (n)	19	19	19	19		19	19	19	19
STD. ERROR (SE)	1.2	0.5	1.6	1.6		0.4	0.3	0.5	0.5
t.05 VALUE FOR n (t.05)	2.101	2.101	2.101	2.101		2.101	2.101	2.101	2.101
(SE)*(t.05) = t.05SE	2.54	0.98	3.41	3.41		0.90	0.59	1.10	1.10
UPPER 95% CONF. INTVL. FOR MEAN	6.1	2.2	8.2	8.2		1.6	1.1	2.3	2.3
LOWER 95% CONF. INTVL. FOR MEAN	1.0	0.3	1.4	1.4		-0.2	-0.1	0.1	0.1
CONVERS. FACTOR ACRES TO SQ. METERS	4046.9	4046.9	4046.9	4046.9					
ESTIMATED NO. OYSTERS IN 1 BUSHEL	1000	500							
ROCK ACREAGE	165	165							
MEAN NO. BUSHELS PER ACRE	14	10	24	24					
UPPER 95% CONF. INTVL. FOR THIS MEAN	25	18	43	43					
LOWER 95% CONF. INTVL. FOR THIS MEAN	4	2	6	6					
MEAN TOTAL BUSHELS IN WHOLE ROCK	2355	1687	4042	4042					
UPPER 95% CONF. INTVL. FOR THIS MEAN	4052	2989	7042	7042					
LOWER 95% CONF. INTVL. FOR THIS MEAN	657	385	1042	1042					

James River Patent Tong Survey

Comparison of Sample Size to Standard Error & Mean

Jail Island - Upper



14

STATISTICAL SUMMARY FOR UPPER JAIL ISLAND (14)

	LIVE OYSTERS				BOXES				SHELL VOL (I)
	SPAT	SMALL	MARKET	TOTAL SML+MKT	SPAT	SML	MKT	TOTAL SML+MKT	
LOW VALUE (PER SQ. METER)	0	0	0	0		0	0	0	0
HIGH VALUE (PER SQ. METER)	56	84		140		20	21	41	9
MEAN NO. PER SQ. METER	5.5	5.6		11.0		1.6	1.5	3.1	1.6
STD.DEV.	9.9	12.8		22.5		3.2	3.0	5.8	1.7
NO. SAMPLES (n)	65	65		65		65	65	65	65
STD. ERROR (SE)	1.2	1.6		2.8		0.4	0.4	0.7	0.2
t.05 VALUE FOR n (t.05)	1.997	1.997		1.997		2.00	2.00	1.997	1.997
(SE)*(t.05) = t.05SE	2.45	3.18		5.57		0.79	0.73	1.44	0.43
UPPER 95% CONF. INTVL. FOR MEAN	7.9	8.8		16.6		2.3	2.2	4.5	2.0
LOWER 95% CONF. INTVL. FOR MEAN	3.0	2.4		5.5		0.8	0.8	1.6	1.1
CONVERS. FACTOR ACRES TO SQ. METERS	4046.9	4046.9							
ESTIMATED NO. OYSTERS IN 1 BUSHEL	1000	500							
ROCK ACREAGE	611.8	611.8							
MEAN NO. BUSHEL PER ACRE	22	45		67					
UPPER 95% CONF. INTVL. FOR THIS MEAN	32	71		103					
LOWER 95% CONF. INTVL. FOR THIS MEAN	12	19		32					
MEAN TOTAL BUSHEL IN WHOLE ROCK	13560	27578		41138					
UPPER 95% CONF. INTVL. FOR THIS MEAN	19624	43337		62961					
LOWER 95% CONF. INTVL. FOR THIS MEAN	7497	11818		19315					

James River Patent Tong Survey

Comparison of Sample Size to Standard Error & Mean

Swash & Swash Mud Slough




Figure 19

STATISTICAL SUMMARY FOR SWASH AND SWASH MUD SLOUGH (15)

	LIVE OYSTERS			BOXES			SHELL VOL (l)	
	SPAT	SMALL	MARKET	TOTAL SML+MKT	SPAT	SML		MKT
LOW VALUE (PER SQ. METER)	0	0	0	0		0	0	0
HIGH VALUE (PER SQ. METER)	129	27	143	143		18	7	20
MEAN NO. PER SQ. METER	20.8	5.6	26.4	26.4		2.4	1.1	3.4
STD. DEV.	24.7	6.0	29.6	29.6		3.2	1.4	4.0
NO. SAMPLES (n)	134	134	134	134		134	134	134
STD. ERROR (SE)	2.1	0.5	2.6	2.6		0.3	0.1	0.3
t.05 VALUE FOR n (t.05)	1.96	1.96	1.96	1.96		1.96	1.96	1.96
(SE)*(t.05) = t.05SE	4.18	1.02	5.01	5.01		0.54	0.24	0.67
UPPER 95% CONF. INTVL. FOR MEAN	25.0	6.6	31.4	31.4		2.9	1.3	4.1
LOWER 95% CONF. INTVL. FOR MEAN	16.6	4.5	21.3	21.3		1.8	0.9	2.8
CONVERS. FACTOR ACRES TO SQ. METERS	4046.9	4046.9	4046.9	4046.9				
ESTIMATED NO. OYSTERS IN 1 BUSHEL	1000	500	500	500				
ROCK ACREAGE	1244.9	1244.9	1244.9	1244.9				
MEAN NO. BUSHEL PER ACRE	84	45	129	129				
UPPER 95% CONF. INTVL. FOR THIS MEAN	101	53	154	154				
LOWER 95% CONF. INTVL. FOR THIS MEAN	67	37	104	104				
MEAN TOTAL BUSHEL IN WHOLE ROCK	104703	56092	160796	160796				
UPPER 95% CONF. INTVL. FOR THIS MEAN	125738	66346	192084	192084				
LOWER 95% CONF. INTVL. FOR THIS MEAN	83669	45838	129507	129507				

James River Patent Tong Survey

Comparison of Sample Size to Standard Error & Mean

Swash - Offshore




Figure 20

STATISTICAL SUMMARY FOR OFFSHORE SWASH (16)

	LIVE OYSTERS				BOXES				SHELL VOL (L)
	SPAT	SMALL	MARKET	TOTAL	SPAT	SML	MKT	TOTAL	
			SML+MKT	SML+MKT		SML+MKT	SML+MKT	SML+MKT	
LOW VALUE (PER SQ. METER)	0	0	0	0	0	0	0	0	0
HIGH VALUE (PER SQ. METER)	190	209	283	283	47	14	28	39	39
MEAN NO. PER SQ. METER	24.5	5.7	30.2	30.2	3.7	1.2	4.0	5.6	5.6
STD. DEV.	44.6	25.8	58.2	58.2	7.0	2.5	6.5	8.4	8.4
NO. SAMPLES (n)	67	67	67	67	67	67	67	67	67
STD. ERROR (SE)	5.5	3.1	7.1	7.1	0.9	0.3	0.8	1.0	1.0
t.05 VALUE FOR n (t.05)	1.992	1.992	1.992	1.992	1.992	1.992	1.992	1.992	1.992
(SE)*(t.05) = t.05SE	10.86	6.27	14.16	14.16	1.71	0.62	1.58	2.05	2.05
UPPER 95% CONF. INTVL. FOR MEAN	35.4	12.0	44.4	44.4	5.4	1.8	5.6	7.6	7.6
LOWER 95% CONF. INTVL. FOR MEAN	13.7	-0.6	16.1	16.1	1.9	0.6	2.4	3.5	3.5
CONVERS. FACTOR ACRES TO SQ. METERS	4046.9	4046.9							
ESTIMATED NO. OYSTERS IN 1 BUSHEL	1000	500							
ROCK ACREAGE	626.51	626.51							
MEAN NO. BUSHEL PER ACRE	99	46	145	145					
UPPER 95% CONF. INTVL. FOR THIS MEAN	143	97	240	240					
LOWER 95% CONF. INTVL. FOR THIS MEAN	55	0	55	55					
MEAN TOTAL BUSHEL IN WHOLE ROCK	62175	28911	91086	91086					
UPPER 95% CONF. INTVL. FOR THIS MEAN	89705	60710	150415	150415					
LOWER 95% CONF. INTVL. FOR THIS MEAN	34644	0	34644	34644					

James River Patent Tong Survey

Comparison of Sample Size to Standard Error & Mean

Jail Island - Lower




Figure 21

STATISTICAL SUMMARY FOR LOWER JAIL ISLAND (17)

	LIVE OYSTERS				BOXES				SHELL VOL(I)
	SPAT	SMALL	MARKET	TOTAL SML+MKT	SPAT	SML	MKT	TOTAL SML+MKT	
LOW VALUE (PER SQ. METER)	0	0	0	0		0	0	0	0
HIGH VALUE (PER SQ. METER)	158	32	183	183		30	7	30	16
MEAN NO. PER SQ. METER	9.3	4.9	14.2	14.2		2.5	1.1	3.6	2.6
STD.DEV.	23.4	7.2	29.2	29.2		5.1	1.6	5.7	3.2
NO. SAMPLES (n)	69	69	69	69		69	69	69	69
STD. ERROR (SE)	2.8	0.9	3.5	3.5		0.6	0.2	0.7	0.4
t.05 VALUE FOR n (t.05)	1.994	1.994	1.994	1.994		1.994	1.994	1.994	1.994
(SE)*(t.05) = t.05SE	5.61	1.74	7.01	7.01		1.23	0.38	1.37	0.78
UPPER 95% CONF. INTVL. FOR MEAN	14.9	6.6	21.2	21.2		3.7	1.4	4.9	3.3
LOWER 95% CONF. INTVL FOR MEAN	3.6	3.2	7.1	7.1		1.3	0.7	2.2	1.8
CONVERS. FACTOR ACRES TO SQ. METERS	4046.9	4046.9							
ESTIMATED NO. OYSTERS IN 1 BUSHEL	1000	500							
ROCK ACREAGE	628.93	628.93							
MEAN NO. BUSHEL PER ACRE	37	40		77					
UPPER 95% CONF. INTVL. FOR THIS MEAN	60	54		114					
LOWER 95% CONF. INTVL. FOR THIS MEAN	15	26		40					
MEAN TOTAL BUSHEL IN WHOLE ROCK	23571	24936		48507					
UPPER 95% CONF. INTVL. FOR THIS MEAN	37856	33768		71623					
LOWER 95% CONF. INTVL. FOR THIS MEAN	9286	16104		25390					

James River Patent Tong Survey

Comparison of Sample Size to Standard Error & Mean

Jail Island - Offshore




Figure 22


STATISTICAL SUMMARY FOR OFFSHORE JAIL ISLAND (18)

	LIVE OYSTERS				BOXES				SHELL VOL (I)
	SPAT	SMALL	MARKET	TOTAL SML+MKT	SPAT	SML	MKT	TOTAL SML+MKT	
LOW VALUE (PER SQ. METER)		0	0	0		0	0	0	0
HIGH VALUE (PER SQ. METER)		83	17	100		25	13	34	30
MEAN NO. PER SQ. METER		7.8	2.4	10.2		4.2	1.0	5.2	4.9
STD. DEV.		11.9	3.4	14.3		5.4	2.0	6.5	5.7
NO. SAMPLES (n)		105	105	105		105	105	105	105
STD. ERROR (SE)		1.2	0.3	1.4		0.5	0.2	0.6	0.6
t.05 VALUE FOR n (t.05)		2.275	2.275	2.275		2.275	2.275	2.275	2.275
(SE)*(t.05) = t.05SE		2.63	0.76	3.17		1.19	0.44	1.45	1.26
UPPER 95% CONF. INTVL. FOR MEAN		10.4	3.2	13.4		5.4	1.5	6.7	6.2
LOWER 95% CONF. INTVL. FOR MEAN		5.2	1.7	7.1		3.0	0.6	3.8	3.6
CONVERS. FACTOR ACRES TO SQ. METERS		4046.9	4046.9						
ESTIMATED NO. OYSTERS IN 1 BUSHEL		1000	500						
ROCK ACREAGE		1017.2	1017.2	1017.2					
MEAN NO. BUSHEL PER ACRE		32	20	51					
UPPER 95% CONF. INTVL. FOR THIS MEAN		42	26	68					
LOWER 95% CONF. INTVL. FOR THIS MEAN		21	14	35					
MEAN TOTAL BUSHEL IN WHOLE ROCK		32109	20151	52260					
UPPER 95% CONF. INTVL. FOR THIS MEAN		42942	26437	69379					
LOWER 95% CONF. INTVL. FOR THIS MEAN		21275	13865	35141					

James River Patent Tong Survey

Comparison of Sample Size to Standard Error & Mean

Wreck Shoal



STATISTICAL SUMMARY FOR WRECK SHOAL (19)

	LIVE OYSTERS				BOXES				SHELL VOL (I)
	SPAT	SMALL	MARKET	TOTAL SML+MKT	SPAT	SML	MKT	TOTAL SML+MKT	
LOW VALUE (PER SQ. METER)	0	0	0	0		0	0	0	0
HIGH VALUE (PER SQ. METER)	76	22	94			20	7	27	28
MEAN NO. PER SQ. METER	6.4	2.3	8.7			4.4	1.0	5.4	9.8
STD.DEV.	12.4	3.9	15.5			4.4	1.5	5.0	7.3
NO. SAMPLES (n)	55	55	55			55	55	55	55
STD. ERROR (SE)	1.7	0.5	2.1			0.6	0.2	0.7	1.0
t.05 VALUE FOR n (t.05)	2.004	2.004	2.004			2.004	2.004	2.004	2.004
(SE)*(t.05) = t.05SE	3.35	1.07	4.20			1.19	0.39	1.34	1.98
UPPER 95% CONF. INTVL. FOR MEAN	9.8	3.3	12.9			5.6	1.4	6.7	11.8
LOWER 95% CONF. INTVL. FOR MEAN	3.1	1.2	4.5			3.2	0.6	4.0	7.9
CONVERS. FACTOR ACRES TO SQ. METERS	4046.9	4046.9							
ESTIMATED NO. OYSTERS IN 1 BUSHEL	1000	500							
ROCK ACREAGE	584.76	584.76							
MEAN NO. BUSHELS PER ACRE	26	18	44						
UPPER 95% CONF. INTVL. FOR THIS MEAN	40	27	66						
LOWER 95% CONF. INTVL. FOR THIS MEAN	12	10	22						
MEAN TOTAL BUSHELS IN WHOLE ROCK	15188	10671	25859						
UPPER 95% CONF. INTVL. FOR THIS MEAN	23107	15716	38823						
LOWER 95% CONF. INTVL. FOR THIS MEAN	7269	5625	12895						

General summary of population sizes

A summary of standing stock estimates for the James River is given in Table 1. The content of this table are distilled from the statistical summaries accompanying Figures 3 through 23. The important conversion figures to acknowledge are that of numbers of small and market oysters per bushel at 1000 and 500 respectively. These correspond to below and above two and one half inches (62.5mm) height (maximum dimension). These summaries do not include young of the year (also commonly termed spat) oysters which are very small and occupy a comparatively negligible volume.

Absolute densities of oysters are highly variable, from high values of 350, 272, 271, 222, 173 and 129 per sq. meter at Upper Horsehead, Lower Horsehead, Moon Rock, Middle Horsehead, V Rock and Point of Shoals respectively, to low values of 14, 11, 10, 9, and 5 at Lower Jail Island, Upper Jail Island, Offshoe Jail Island, Wreck Shoal, and Swash respectively. Mean estimates of standing stocks of seed (small) and market oysters are 465,356 and 258,869 bushels respectively, for a total of approximately 724,225 bushels in the surveyed section of James River. The confidence interval around these values gives upper and lower values of 318,542 and 612,169 bushels for seed (small), and 155,582 and 365,078 bushels for market oysters respectively. A limited number of individual rocks had lower estimates of zero for market oysters - these reflect analysis of data that include a large number of samples with zero market size oysters present.

Substantial seed (small) oyster resources are present in a number of locations: Upper Deep Water Shoal, the components of Horsehead Rock, V Rock, Point of Shoals, Cross Rock, and the large areas of Swash and Jail Island. The bulk of market oysters are located on the same rocks.

In the Rappahannock River standing stock estimates were made for Carters Rock, Ross's Rock, Bowlers Rock, Long Rock, and Sharps Rock (inshore). These are all very small rocks and of limited commercial importance. The estimated seed oysters resources on these were 126, 637, 36, 78, and 13 bushels respectively. The estimated market oyster resources on these were 69, 371, 79, 202 and 0 bushels respectively. Only Ross's Rock supported any commercial activity in the public oyster season of 1993-94.

Table 1

JAMES RIVER OYSTER ROCKS
 Standing Stock Estimates
 BUSHELS OF OYSTERS IN ROCK ACREAGE SAMPLED
 Fall 1993

Chart No.	OYSTER ROCK Name	ACRES SAMPLED	SEED OYSTERS			MARKET OYSTERS			SEED & MARKET COMBINED		
			MEAN ESTIMATE	Lower Mean Estimate	Upper Mean Estimate	MEAN ESTIMATE	Lower Mean Estimate	Upper Mean Estimate	MEAN ESTIMATE	Lower Mean Estimate	Upper Mean Estimate
1	Upper Deep Water Shoal	233.92	46,472	31,692	61,252	37,579	25,918	49,240	84,051	57,610	110,492
2	Lower Deep Water Shoal	19.93	798	508	1,089	1,371	969	1,773	2,170	1,477	2,862
3	Upper Horsehead	3.009	3,588	429	6,747	1,348	101	2,594	4,936	530	9,341
4	Middle Horsehead	19.465	16,877	11,671	22,083	1,158	521	1,795	18,035	12,192	23,878
5	Lower Horsehead	19.467	19,963	15,439	24,487	2,954	1,179	4,729	22,917	16,619	29,216
6	Moon Rock	3.95	3,948	1,744	6,152	791	223	1,360	4,740	1,967	7,512
7	V Rock	72.053	45,950	39,780	52,120	11,842	9,777	13,906	57,792	49,557	66,026
8	Point of Shoals	131.71	55,906	42,919	68,893	25,463	19,309	31,617	81,369	62,229	100,509
9	Cross Rock	36.688	11,151	7,050	15,252	2,329	1,685	2,972	13,480	8,735	18,224
10	Shanty Rock	3.58	471	32	909	65	0	160	536	32	1,069
11	Dry Lumps	5.93	360	142	578	40	0	90	400	142	667
12	Mulberry Point	86.85	6,436	2,973	9,899	3,937	2,240	5,633	10,372	5,213	15,532
13	Swash	165	2,355	657	4,052	1,687	385	2,989	4,042	1,042	7,042
14	Upper Jail Island	611.8	13,560	7,497	19,624	27,578	11,818	43,337	41,138	19,315	62,961
15	Swash-Swash Mud Slough	1244.9	104,703	83,669	125,738	56,092	45,838	66,346	160,795	128,507	192,084
16	Offshore Swash	626.51	62,175	34,644	89,705	28,911	0	60,710	91,086	34,644	150,415
17	Lower Jail Island	628.93	23,571	9,286	37,856	24,936	16,104	33,768	48,507	25,390	71,623
18	Offshore Jail Island	1017.2	31,884	21,141	42,626	20,117	13,890	26,343	52,090	35,030	68,970
19	Wreck Shoal	584.76	15,188	7,269	23,107	10,671	5,625	15,716	25,859	12,895	38,823
TOTALS		5515.652	465,356	318,542	612,169	258,869	155,582	365,078	724,225	474,124	977,247

Size distribution data and implications for interpretation of general summary

Size distribution data, by numbers of individual oysters present within each 5 mm height size class interval, for all 19 areas sampled is illustrated in Figure 24. For convenience this is displayed as six graphics. Again, young of the year (spat) oysters are not included in this illustration. The dominant feature of all plots is the rapid decrease in number of individuals in all locations above the 60-65 mm (mid point 63 mm on Figure 24) size class. This corresponds closely with the two and one half inch (62.5 mm) minimum size for market oyster harvest, suggesting efficient harvesting above the size limit. Despite this individual of over 100 mm maximum dimension were found in very limited numbers in the majority of locations. The size distribution data illustrate that the increase in minimum size for market oyster exploitation to three inches (76 mm) for the 1994-1995 public oyster season may result in some hardship to watermen in that large numbers of individuals were at or below 60 mm in the Fall of 1993, and would have to grow substantially through the spring and summer of 1994 to attain a 76 mm size and become available to the fishery in the Fall of 1994.

When size distribution data by individual numbers is replotted by either biomass or potential contribution to egg production other facets of stock management are implicated. A sample of 111 oysters of sizes varying from 30 to 95 mm height was collected from the James River on May 13, 1994 and used to generate conversion functions of height to live weight and dry tissue weight. The relationships generated by MINITAB analysis are:

$$\text{live weight (g)} = 0.0064642 \times \text{height}^{2.1095}$$

$$\text{dry tissue weight (g)} = 0.000423 \times \text{height}^{1.7475}$$

In turn these can be related to fecundity by the relationship:

$$\text{fecundity (millions of eggs)} = 39.06 \times \text{dry tissue weight}^{2.36}$$

These conversions are illustrated for data from Lower and Upper Deep Water Shoal in Figure 25. In the instance of both live and dry tissue weight the mode moves above the 60-65 mm size class, illustrating the importance of the numerically smaller size classes in the ecologically important processes of filtration and benthic-pelagic coupling. The fecundity issue is critical, in that the basis for setting minimum harvest size is to maximize reproductive output prior to harvest (although this is somewhat questionable in the James River where, until the 1994-1995 season, seed harvest procedures allowed removal of essentially all oysters from the majority of public oyster ground). When considering contribution to egg production 76 and 65% of production for Lower and Upper Horsehead is in the 60-65 mm size class and above. Harvesting these size classes, despite their numerical inferiority to smaller size classes, can clearly have major impact on egg production. Note that these percentages are calculated giving equal weighting to sex ration with increasing size. Although a matter of debate in the literature the positions vary from unity of ratio with size to a predominance of females with larger size classes. If the latter were the case then the 76 and 65% values are conservative estimates!. An increase of minimum size to 76 mm decreases these values considerably: 48 and 32% of estimated egg production comes from individuals in the 75-80 mm size interval and larger in the two locations. Increasing the minimum size limit for market oysters from two and one half to three inches (62.5 to 76 mm) effectively doubles the available egg production from the resource.

Figure 24: Size class distribution




Figure 24: Size class distribution




Figure 24: Size class distribution




Figure 24: Size class distribution




Figure 24: Size class distribution




Figure 24: Size class distribution





Figure 25: Comparison of biomass distribution by live and dry tissue weights, and egg production by size class interval



Fishery Dependent Methods

Barber and Mann (1991), supported by Chesapeake Bay Stock Assessment Committee (CBSAC) funds, employed Leslie - DeLury analysis of commercial fishery data (daily and weekly boat count data to estimate effort, landing data to estimate catch, both data sets collected and provided by the VMRC) to estimate recent decline in standing stock of oysters in the James River. A secondary objective of the current study was to compare, where possible, fishery independent and fishery dependent estimates of standing stock.

Fishery-dependent estimates quantify only that portion of the resource harvested as a part of commercial activity and systematically exclude smaller animals that also contribute to gamete production. They are also susceptible to errors from under-reporting of catch. An excellent review of the limitations of Leslie-DeLury application to invertebrate fisheries is given by Breen (1992). Leslie-DeLury analysis becomes less sensitive as stocks become very low or if there is not a marked decrease in catch per unit effort over a fishing season. Both situations are likely to occur in application to the Virginia oyster fishery, and are complicated by variable, low or no effort towards the end of the fishing season as watermen turn to other resources. Finally, any method based on regression analysis must comply to certain statistical prerequisites, including normal distribution of data. This may not always be the case in data obtained from commercial fishing operations. Despite these limitations, fishery dependent estimates of standing stock have been the only estimates available to the Virginia Marine Resource Commission.

Tables 3 and 4 provide summaries, taken from V.M.R. C. records, of seed and market oyster catches, by month, for the public fishery for the seasons from 1982-1983 through 1994-1994. To illustrate trends in seasonal totals the values are plotted in Figure 26. The period of 1982-1983 through 1988-1989 are characterized by an initial focus on the market oyster fishery, with an accompanying decline, subsequently followed by an increasing harvest of seed oysters in the 1986-1987 through 1988-1989 period. Note that the focus on market oysters is historically unprecedented in this location. Previously sub sets of the data of Tables 3 and 4 were used to generate Leslie-DeLury estimates of standing stock of market oysters, and the data given limited distribution (including the proposal which lead to the current fishery independent assessment). These estimates are given in Table 5. They suggest a rapidly diminishing resource, and values that are well below those suggested by the fishery independent estimates given earlier in this report. The immediate question to present is why? The answer to this question is that the analysis is probably flawed!

Leslie-DeLury analysis relies on decreases in catch per unit effort over time to estimate initial values of standing stock. The data as used for effort do not distinguish between effort devoted to market oyster harvest and that devoted to seed oyster harvest. On any particular boat at any time in the period described by Table 5 attention may have been devoted to market oysters or seed oysters in isolation, or to the peculiar (to this River, and again a product of the regulations allowing both seed and market oyster harvest from the same location) activity of "two piling" - retaining both seed and market oysters as separate catches for inspection purposes on the same vessel. Short term variability in relative effort devoted to each resource, suggested by catch landing data in Tables 3 and 4, will complicate the estimation of effort. Although "two piling" was not a common practice (market oyster prices dictating more efficient use of limited space) such activity also influenced effort data on each resource. To rectify the problem and repeat the analysis would probably require that all oyster tax records provided from individual harvesters be re-examined and a new database generated which consistently distinguishes, if possible, between the two catches. In the interim period the continued use of summary data that does not distinguish effort between the two resources should not be used to generate fishery independent estimates of standing stock.

Table 2: James River Seed and Market Oyster Harvest: 1982-1983 through 1993-1994.

HARVEST SEASON
SEED OYSTERS

	OCTOBER	NOVEMBER	DECEMBER	JANUARY	FEBRUARY	MARCH	APRIL	MAY	JUNE	TOTAL
82/83	83,282	72,595	57,144	29,264	24,245	47,418	51,110	56,599	23,536	445,193
83/84	68,774	68,414	32,939	15,043	35,605	15,947	53,375	33,961	22,076	346,134
84/85	72,463	54,338	38,744	25,277	28,649	50,801	71,595	54,119	13,881	409,867
85/86	61,420	48,731	32,908	32,240	16,920	36,238	27,745	20,301	0	276,503
86/87	39,092	34,193	14,619	3,839	3,375	18,215	41,088	47,985	0	202,406
87/88	20,592	16,281	11,634	1,017	17,585	29,548	15,949	21,847	0	134,453
88/89	13,827	1,948	3,934	675	797	1,281	4,757	14,084	0	41,303
89/90	7,523	8,817	3,137	4,915	6,062	6,266	7,633	7,031	0	51,383
90/91	624	1,475	2,476	4,670	16,130	8,539	16,476	4,620	0	55,010
91/92	1,996	1,320	3,842	17,109	16,052	10,020	2,011	1,187	0	53,537
92/93	1,896	757	3,135	18,927	18,142	38,824	925			76,606
93/94	31917	49897	38870	7810	3060	5150	4847			141551

MARKET OYSTERS

	OCTOBER	NOVEMBER	DECEMBER	JANUARY	FEBRUARY	MARCH	APRIL	MAY	JUNE	TOTAL
82/83	2,039	4,684	2,384	2,004	1,182	386	1,003	1,493	956	16,131
83/84	6,833	8,339	3,423	9,492	3,423	3,420	6,018	6,171	1,632	48,746
84/85	370	6,912	1,694	1,904	1,259	1,243	2,452	3,087	1,555	21,467
85/86	2,196	706	0	4,516	3,893	3,882	3,145	10,418	0	28,756
86/87	62,675	62,212	70,346	50,139	52,823	21,958	15,867	6,764	0	342,784
87/88	65,268	57,052	46,343	36,965	31,433	28,029	17,235	15,449	0	297,774
88/89	43,113	25,225	22,876	15,220	10,997	14,590	7,461	7,474	0	146,956
89/90	11,699	10,658	7,235	13,873	6,587	7,745	6,157	4,588	0	68,542
90/91	7,747	8,126	6,350	4,582	3,487	3,461	4,269	5,384	0	43,406
91/92	4,709	3,651	4,326	2,404	2,373	2,719	3,088	2,314		25,584
92/93	3,584	1,987	2,774	2,161	1,117	2,117	2,153	334		16,227
93/94	1448	2464	2077	403	135	285	236			7048

Table 3: James River Boat Days (Effort) and Catch / Day: 1982-1983 through 1993-1994


HARVEST SEASON BOAT DAYS	OCTOBER	NOVEMBER	DECEMBER	JANUARY	FEBRUARY	MARCH	APRIL	MAY	JUNE	TOTAL
82/83	1,142	1,225	895	704	564	692	769	834	262	7,087
83/84	1,260	1,305	674	710	809	646	961	786	382	7,533
84/85	1,392	1,045	1,112	331	532	866	975	1,024	260	7,537
85/86	1,078	731	668	626	557	636	730	599	0	5,625
86/87	2,408	2,518	2,752	1,966	2,222	1,158	1,432	1,298	0	15,754
87/88	3,628	3,021	2,939	2,216	3,081	3,042	1,423	1,775	0	21,305
88/89	3,355	2,139	2,360	1,554	1,193	1,340	1,099	987	0	14,027
89/90	1,758	1,362	1,019	1,840	1,075	1,154	899	703	0	9,810
90/91	892	967	1,003	831	825	578	828	774	0	6,698
91/92	682	488	500	600	534	561	458	176	0	4,032
92/93	294	291	274	306	333	833	165	202	0	2,698

CATCH/DAY	OCTOBER	NOVEMBER	DECEMBER	JANUARY	FEBRUARY	MARCH	APRIL	MAY	JUNE	TOTAL
82/83	75	63	67	44	45	69	68	70	94	65
83/84	60	59	54	35	48	30	62	51	62	52
84/85	52	59	37	82	56	60	76	56	59	57
85/86	59	68	49	59	37	63	42	51	0	54
86/87	42	38	31	28	25	35	40	42	0	35
87/88	24	24	20	17	16	19	23	21	0	20
88/89	17	13	11	10	10	12	11	22	0	13
89/90	11	14	10	10	12	12	15	17	0	12
90/91	9	10	9	11	23	21	25	13	0	15
91/92	10	10	11	31	34	20	11	20	0	20
92/93	19	9	22	69	58	42	19	20	0	34

Table 5. Previously generated estimates of standing stock of oysters (bushels) in James River, estimated using Leslie-DeLury from partial (October-December) and full (October-May) season commercial fishery data. The notation ns indicates no significant estimate available from regression analysis. We now suggest that this analysis is compromised by the lack of separation of estimates of effort in the seed and market oyster fisheries, and that these data should be viewed with caution.

Year	October-December	October -May
1986-1987	1,093,000	618,000
1987-1988	840,500	ns
1988-1989	429,000	ns
1989-1990	234,000	265,750
1990-1991	78,400	95,350

**FIGURE 26: OYSTER PRODUCTION BY SEASON:
1982/3 THROUGH 1993/4**



SEASON	SEED OYSTERS	MARKET OYSTERS
82/83	45,000	445,193
83/84	48,746	346,134
84/85	21,467	409,867
85/86	28,756	276,503
86/87	202,406	342,784
87/88	134,453	297,774
88/89	41,303	146,956
89/90	51,383	68,542
90/91	55,010	43,406
91/92	53,537	25,584
92/93	76,606	16,227
93/94	141,551	7,048

Literature cited

- Barber, B. J. and R. Mann. 1991. Estimation of Standing Stock of Oysters in the James River, Virginia, Using Commercial Fishing Records. Virginia Institute of Marine Science, Special Report in Applied Marine Science and Ocean Engineering, No 310.
- Breen, P. A. 1992. A review of models used for stock assessment in abalone fisheries. pp 253-275 in: Abalone of the World: Biology, Fisheries and Culture. S. A. Shepherd, M. J. Tegner, S. A. Guzman del Proo (Eds.). Fishing News Books, Blackwell Scientific, 608p.
- Bros, W. E. and B. C. Cowell. 1987. A technique for optimizing sample size (replication). J. Exp. Mar. Biol. Ecol. 114: 63-71.
- Chai, A-L, M. Homer, C-F. Tsai and P. Gouletquer. 1992. Evaluation of oyster sampling efficiency of patent tongs and an oyster dredge. North American Journal of Fisheries Management 12: 825-832.
- Cox, C. 1988. Seasonal changes in the fecundity of oysters Crassostrea virginica (Gmelin) from four oyster reefs in the James River, Virginia. M.A. thesis, College of William and Mary, Williamsburg, VA.
- Cox, C. and R. Mann. 1992. Fecundity of oysters, Crassostrea virginica (Gmelin), in the James River, Virginia, U.S.A. J. Shellfish Res. 11(1): 47-52.
- Hargis, W. J. Jr. and D. S. Haven. 1988. The imperilled oyster industry of Virginia. VIMS Special Report Number 290 in Applied Marine Science and Ocean Engineering. 130p.
- Haven, D. S., W. J. Hargis, Jr. and P. Kendall. 1981. The present and potential productivity of the Baylor Grounds in Virginia. Va. Inst. Mar. Sci., Spec. Rep. Appl. Mar. Sci. Ocean. Eng. No. 243: 1-154.
- Haven, D. S. and J. P. Whitcomb. 1983. The Origin and Extent of Oyster Reefs in the James River, Virginia. J. Shellfish Res. 3(2):141-151
- Haven, D. S. and J. P. Whitcomb. 1989. The Location and Topography of oyster reefs in the Rappahannock River Estuary, Virginia. J. Shellfish Res. 8:105-116
- Morales-Alamo, R. and R. Mann. 1990. Recruitment and growth of oysters on shell planted at four monthly intervals in the Lower Potomac River, Maryland. J. Shellfish Res. 9(1): 165-172).
- Rheinhardt, R. D. and R. Mann. 1990. Development of epibenthic fouling communities on a natural oyster bed in the James River, Virginia. Biofouling: 2: 13-25.

Conclusions and Recommendations

The James River will remain the only substantial source of both seed and market oysters for the public fishery for the immediate future. The resource in the Rappahannock will remain of minor importance to the total fishery production. The James River market oyster harvest for the 1993-94 public season of 5,173 bushels represents approximately 2.2% of the estimated standing stock; however, the seed harvest of 72,470 bushels for the same period represents approximately 15.6% of the estimated standing stock. These are the first ever fishery independent estimates, so long term comparisons of harvest versus standing stock are not possible, although such levels of exploitation appear reasonable at this time. There is considerable spatial variability in oyster density, and harvesting will probably continue to focus on those areas with high density such as Horsehead, Moon Rock, V Rock and Point of Shoals. The seed resource is still substantial, but its utilization will probably be controlled by factors other than availability to the watermen. Lease holding planters are reluctant to purchase seed oysters that may have already been exposed to disease. While mortalities associated with disease are limited in the upper part of the James as sampled oysters may contract infective particles. When transferred to higher salinity grow out sites infected oysters essentially participate in a race between the progressing disease and growth to market size. The financial consequences to the planter of disease related loss in this instance has prompted caution in seed sales. Seed prices, when buyers are present, are variable, but often suppressed.

The recent increase in minimum size of market oysters may suppress landings in the 1994-1995 season depending on the growth of smaller size classes in the spring and summer of 1994. The majority of oysters were below the former size limit during the Fall 1993 sampling, and consistently good growth would be required to make them abundant for the 1994-1995 public season. From an ecological perspective the increase in minimum size is to be applauded. Calculations based upon observed size class distribution data illustrate the important contribution of the numerically limited market oysters to total biomass, and hence benthic pelagic coupling. More importantly, accompanying calculations suggest that the modest increase in minimum size will double the available egg production from remaining oysters - a clear bonus in a long term plan to rebuild the resource.

The nature of fishery dependent data records is such that they do not adequately distinguish between market and seed oyster fishing activity, and changes in emphasis from one to the other cause major variability in catch per unit effort data. In turn this compromises the value of standing stock estimates obtained by Leslie-DeLury analysis. We suggest they not be used until fishery records can be re-examined for possible generation of a new database. Fishery independent estimates appear to provide a much more stable method of stock estimate for management purposes.